



## RESEARCH ARTICLE

### Hematological effects of Coenzyme Q10 in streptozotocin-induced diabetic rats

Ercan Keskin<sup>1</sup>, Deniz Uluşık<sup>1\*</sup>

<sup>1</sup>Selçuk Üniversitesi, Veteriner Fakültesi, Fizyoloji Anabilim Dalı, Konya, Türkiye

Received: 13.02.2017, Accepted: 05.04.2017

\* denizfedai@selcuk.edu.tr

### Streptozotosin ile diyabet oluşturulan ratlarda Koenzim Q10 (CoQ10)'un hematolojik etkileri

Eurasian J Vet Sci, 2017, 33, 3, 167-171

DOI:10.15312/EurasianJVetSci.2017.154

#### Öz

**Amaç:** Bu çalışmada streptozotosin ile diyabet oluşturulan ratlarda Koenzim Q10 (CoQ10)'un bazı hematolojik parametreler üzerine etkilerinin belirlenmesi amaçlandı.

**Gereç ve Yöntem:** Bu amaçla 38 adet sağlıklı, yetişkin erkek rat beş gruba ayrıldı. Dört hafta boyunca 1. gruptaki hayvanlar standart rat yemi ile beslenirken, 2. gruptaki hayvanlara dört hafta boyunca günlük 0,3 ml mısır yağı intraperitoneal (IP) olarak uygulandı. Aynı süre ile 3. gruptaki hayvanlara günlük 10 mg/kg CoQ10 IP olarak enjekte edildi. 4. gruptaki hayvanlarda 40 mg/kg streptozotosinin günde tek doz olmak üzere iki gün subkutan (SC) enjeksiyonu ile diyabet oluşturulurken, aynı yolla diyabet oluşturulan 5. gruptaki hayvanlara dört hafta boyunca günlük 10 mg/kg CoQ10 IP olarak enjekte edildi. Bütün hayvanlardan alınan kan örneklerinde alyuvar (RBC) ve akyuvar (WBC) sayıları, hemoglobin miktarı, hematokrit değeri (PCV), lökosit yüzdeleri ile ortalama alyuvar hacmi (MCV), ortalama alyuvar hemoglobini (MCH) ve ortalama alyuvar hemoglobin konsantrasyonu (MCHC) belirlendi.

**Bulgular:** Streptozotosin ile diyabet oluşturulan ratların RBC, hemoglobin miktarı ve hematokrit düzeylerinin kontrol grubuna göre anlamlı olarak düşük olduğu ( $p<0.05$ ) belirlendi. Diyabetli hayvanlara CoQ10 uygulaması ile hematokrit değeri diyabetli grupla karşılaştırıldığında önemli artış ( $p<0.05$ ) gösterdi. DeneySEL diyabet WBC ve nötrofil yüzdesinde kontrole göre önemli artışlara ( $p<0.05$ ), lenfosit yüzdesinde ise azalmaya ( $p<0.05$ ) neden oldu. Diyabetli hayvanlara CoQ10 uygulaması ile diyabetli gruba göre nötrofil yüzdesinde önemli azalma ve lenfosit yüzdesinde ise önemli artış belirlendi ( $p<0.05$ ).

**Öneri:** Elde edilen bu sonuçlar diyabetli ratlara CoQ10 uygulamasının diyabetik hasara bağlı hematolojik parametrelerdeki olumsuz değişiklikler üzerine yararlı etkilere sahip olabileceğini göstermiştir.

**Anahtar kelimeler:** CoQ10, diyabet, hematolojik parametreler, rat

#### Abstract

**Aim:** This study was conducted to evaluate the effects of Coenzyme Q10 on some hematological parameters in streptozotocin-induced diabetic rats.

**Materials and Methods:** In the experiment, 38 healthy, adult male rats were divided into five groups. Group 1 was fed standard rat pellets for four weeks. Group 2 was administered with 0,3 ml corn oil IP daily for four weeks, group 3 was injected IP with 10 mg/kg CoQ10 daily for four weeks, group 4 was made diabetic by SC injections of 40 mg/kg streptozotocin as single daily dose for two days, group 5 was made diabetic by SC injections of streptozotocin in the same way and then was injected IP with 10 mg/kg CoQ10 daily for four weeks. In blood samples received from all animals, red blood cells (RBC) and white blood cells (WBC) counts, hemoglobin amount, hematocrit value (PCV), differential leucocyte counts and mean cell volume (MCV), mean corpuscular hemoglobin (MCH), mean corpuscular hemoglobin concentration (MCHC) were determined.

**Results:** In diabetic rats, RBC, hemoglobin and hematocrit values were determined to be lower ( $p<0.05$ ) than the control group. CoQ10 application to diabetic animals increased ( $p<0.05$ ) hematocrit value compared diabetic group. Experimentally induced diabetes caused significantly ( $p<0.05$ ) increments in WBC count and neutrophil percentage, while lymphocyte percentage was decreased ( $p<0.05$ ) compared to control group. CoQ10 treatment to the diabetic group markedly ( $p<0.05$ ) decreased the neutrophil percentage and increased the lymphocyte percentage compared to diabetic group.

**Conclusion:** These results showed that CoQ10 treatment to diabetic rats may have some beneficial properties on hematological parameters negatively affected from diabetic injury.

**Keywords:** CoQ10, diabetes, hematological parameters, rats





## Introduction

Diabetes mellitus (DM) is a metabolic disorder with different etiologies. It is characterized by insufficiency to regulate blood glucose level caused by relative or absolute deficiency in insulin. Diabetes mellitus may occur as a result of pancreatic  $\beta$ -cells dysfunction, which causes reduction in insulin secretion. The disease could also occur due to the insulin receptors resistant to the functions of circulating insulin (ADA 2010). Repetitive or obstinate hyperglycemia during diabetes causes formation glycated body proteins. This effect leads to secondary complications affecting eyes, kidneys, nerves and arteries (Sharma 1993, Aladodo et al. 2013). Erythrocytes (RBC) are seriously affected by hyperglycemia. It was reported these effects of hyperglycemia such as altered membrane lipid composition, reduced filterability, glycated hemoglobin, and the accumulation of advanced glycosylation end-products on the membrane (Schmid-Schönbein and Volger 1976, Wautier et al. 1994, Labrousche et al. 1996, Manodori and Kuypers 2002). In diabetes, reduced hemoglobin has been reported (Mansi 2006) and it may be associated with the decrease in the red blood cell count and packed cell volume (Moss 1999, Muhammad and Oloyede 2009, Aladodo et al. 2013).

Coenzyme Q10 (CoQ10) has been focused as a dietary supplement capable of influencing cellular bioenergetics and counteracting on some damage caused by free radicals (Linnae et al. 2002, Butler et al. 2003, Rosenfeldt et al. 2003, Zhou et al. 2005). CoQ10 is known as a vitamin like and fat-soluble substance existing in all cells (Crane 2001). It is strictly related with various activities including the transferring of electrons within the mitochondrial oxidative respiratory chain and ATP production. Further, it acts as [1] an essential antioxidant and supporting the regeneration of other antioxidants, [2] affecting the stability, fluidity and permeability of membranes, [3] stimulating cell growth and inhibiting cell death (Niki 1997, Crane 2001, Cooke et al. 2008).

Hence, the objective of the present study was to investigate hematological effects of CoQ10 in streptozotocin-induced diabetic rats.

## Materials and Methods

We used 38 healthy, adult male Wistar Albino rats. The animals were kept in individual cages during the four weeks experiment and allowed free access to water and standard pellets. Diabetes was induced by SC injection of streptozotocin (Sigma-Aldrich, St. Louis, MO, USA) at a dose of 40 mg/kg daily in 0.1 M citrate buffer (pH 4.5) single daily dose for two days. To prevent the streptozotocin-induced hypoglycemia, rats received 5% dextrose solution after 6 h of streptozotocin administration for next 3 days. After 1 week, induction of diabetes was verified by measuring blood glucose level with strips using glucometer (PlusMED Accuro, Taiwan) via the

tail vein. Animals whose blood glucose level higher than 250 mg/dL were considered diabetic and included in the experiments. The mean weights of all groups were similar. The rats were divided into five groups. Group 1 (n=6) was fed standard rat pellets for four weeks, group 2 (n=6) was administered at 0,3 ml corn oil IP daily for four weeks, group 3 (n=6) was injected IP with 10 mg/kg CoQ10 (Sigma-Aldrich, St. Louis, MO, USA) daily for four weeks, group 4 (n=7) was made diabetic by SC injections of streptozotocin at dose of 40 mg/kg in 0.1 M citrate buffer (pH 4.5) single daily dose for two days, group 5 (n=9) was made diabetic by SC injections of streptozotocin in the same way and then was injected IP with 10 mg/kg CoQ10 daily for four weeks. During the experiment, three animals from group 4 and one animal from group 5 were died due to streptozotocin-induced hypoglycemia.

At the end of the study, blood samples were taken from all animals. In these blood samples, red blood cells (RBC) and white blood cells (WBC) counts, hemoglobin amount, hematocrit value (PCV), differential leucocyte counts and mean cell volume (MCV), mean corpuscular hemoglobin (MCH), mean corpuscular hemoglobin concentration (MCHC) were determined using an automated cell counter (Abbott Cell Dyn 3700, Chicago, USA). The Ethical Committee of Selcuk University Experimental Medicine Research and Application Center (Report no. 2015-50) approved the study protocol.

The data were analyzed using one-way ANOVA (SPSS 17). Differences among the groups were determined by Duncan's multiple range test. Differences were considered significant at  $p < 0.05$ .

## Results

In streptozotocin induced diabetic rats, RBC, hemoglobin and hematocrit values were determined to be significantly lower (Table 1,  $p < 0.05$ ) than the control group. There were no any changes in respect to the same parameters with CoQ10 treatment alone compared to the control group. CoQ10 application for four weeks to diabetic animals alleviated the reducing in some hematologic parameters compared with the diabetic group levels. But the changes in hematocrit value was important (Table 1,  $p < 0.05$ ) in the diabetic group with CoQ10 treatment. MCV, MCH and MCHC levels didn't show significant differences in all groups. On the other hand, experimentally induced diabetes caused significant (Table 2,  $p < 0.05$ ) increments in WBC count and neutrophil percentage, while lymphocyte percentage was relatively decreased (Table 2,  $p < 0.05$ ) compared to control group. CoQ10 treatment to the diabetic group markedly (Table 2,  $p < 0.05$ ) decreased the neutrophil percentage and increased relative lymphocyte percentage compared to diabetic group (Table 2,  $p < 0.05$ ). In the same group, WBC count tended to decline but this was not important compared to diabetic group. In oil and CoQ10 group, WBC count and differential leucocyte values weren't

Table 1. Mean RBC, Hb, PCV, MCV, MCH and MCHC values (Mean  $\pm$  SE)

Groups	RBC ( $\times 10^6/\text{mm}^3$ )	Hb (g/dl)	PCV (%)	MCV ( $\mu\text{m}^3$ )	MCH (pg)	MCHC (%)
Group 1	7,40 $\pm$ 0,37 <sup>a</sup>	14,78 $\pm$ 0,63 <sup>a</sup>	41,23 $\pm$ 0,99 <sup>a</sup>	56,46 $\pm$ 3,15	20,06 $\pm$ 0,49	35,95 $\pm$ 1,67
Group 2	7,25 $\pm$ 0,43 <sup>ab</sup>	14,05 $\pm$ 0,59 <sup>ab</sup>	40,57 $\pm$ 1,61 <sup>a</sup>	56,71 $\pm$ 3,21	19,73 $\pm$ 1,37	34,69 $\pm$ 1,02
Group 3	7,61 $\pm$ 0,55 <sup>a</sup>	15,25 $\pm$ 0,40 <sup>a</sup>	42,87 $\pm$ 1,16 <sup>a</sup>	57,48 $\pm$ 3,48	20,46 $\pm$ 1,23	35,65 $\pm$ 0,99
Group 4	6,28 $\pm$ 0,26 <sup>b</sup>	11,71 $\pm$ 0,54 <sup>c</sup>	34,40 $\pm$ 0,96 <sup>b</sup>	55,14 $\pm$ 2,22	18,79 $\pm$ 1,09	34,13 $\pm$ 1,61
Group 5	7,11 $\pm$ 0,14 <sup>ab</sup>	13,07 $\pm$ 0,37 <sup>bc</sup>	39,68 $\pm$ 0,74 <sup>a</sup>	55,97 $\pm$ 1,57	18,47 $\pm$ 0,78	33,03 $\pm$ 1,12

The difference between mean values with different superscripts in the same column is significant for each parameter;  $p < 0.05$ . Group 1, control; group 2, oil; group 3, CoQ10; group 4, diabetes; group 5, CoQ10 and diabetes.

Table 2. Mean WBC and differential leukocyte counts (Mean  $\pm$  SE)

Groups	WBC ( $\times 10^3/\text{mm}^3$ )	Neutrophils (%)	Lymphocytes (%)	Monocytes (%)	Eosinophils (%)	Basophiles (%)
Group 1	9,75 $\pm$ 1,55 <sup>b</sup>	21,43 $\pm$ 2,48 <sup>b</sup>	72,42 $\pm$ 2,26 <sup>a</sup>	4,17 $\pm$ 0,65	1,53 $\pm$ 0,60	0,45 $\pm$ 0,21
Group 2	10,78 $\pm$ 0,43 <sup>ab</sup>	20,25 $\pm$ 0,95 <sup>b</sup>	74,25 $\pm$ 1,75 <sup>a</sup>	3,83 $\pm$ 0,87	1,33 $\pm$ 0,49	0,33 $\pm$ 0,21
Group 3	10,52 $\pm$ 0,70 <sup>ab</sup>	19,77 $\pm$ 0,77 <sup>b</sup>	74,55 $\pm$ 0,96 <sup>a</sup>	3,63 $\pm$ 0,80	1,67 $\pm$ 0,42	0,38 $\pm$ 0,24
Group 4	13,18 $\pm$ 0,98 <sup>a</sup>	31,96 $\pm$ 3,26 <sup>a</sup>	61,16 $\pm$ 3,72 <sup>b</sup>	4,71 $\pm$ 0,87	1,75 $\pm$ 0,29	0,41 $\pm$ 0,20
Group 5	11,13 $\pm$ 0,54 <sup>ab</sup>	25,18 $\pm$ 1,40 <sup>b</sup>	68,40 $\pm$ 1,94 <sup>a</sup>	4,31 $\pm$ 0,53	1,62 $\pm$ 0,37	0,49 $\pm$ 0,17

The difference between mean values with different superscripts in the same column is significant for each parameter;  $p < 0.05$ . Group 1, control; group 2, oil; group 3, CoQ10; group 4, diabetes; group 5, CoQ10 and diabetes.

different from the control group levels.

## Discussion

The reductions in RBC, hemoglobin and hematocrit levels of the diabetic group when compared to the control group were seem to consistent with other studies which used similar experimental diabetes (Mansi 2006, Aladodo et al. 2013). The reasons of these changes are attributed to oxidative stress, decreasing of erythrocyte lifespan, bone marrow suppression. It has been suggested that anemia resulting from diabetes mellitus are due to peroxidation of membrane lipids, decreasing of membrane fluidity, oxidation of glycosylated membrane proteins and hemolysis of RBC (Kennedy and Baynes 1984, Bakan et al. 2006, Mansi 2006).

In this study, CoQ10 treatment to diabetic animals alleviated the reducing in some hematologic parameters compared with the diabetic group levels. The changes in hematocrit value was important (Table 1,  $p < 0.05$ ) in the diabetic group treated with CoQ10. It has been well known that elevated glucose causes oxidative stress as a result of increased production of reactive oxygen species (ROS), nonenzymatic glycation of proteins and glucose autooxidation in diabetes mellitus (Brownlee 2001, Modi et al. 2006). The positive effect of CoQ10 on hematocrit value of diabetic animals could

be based on its some antioxidant properties. CoQ10 is recognized as a powerful systemic radical scavenger (Prosek et al. 2008). Moreover, there are many findings related to antioxidant effects of CoQ10 in diabetes mellitus. Niklowitz et al. (2004) found that CoQ10 is greater antioxidant than Vitamin E. In another research, it has been suggested that CoQ10 enhanced the availability of other antioxidants such as Vitamin E, Vitamin C and beta-carotene (Shekelle et al. 2003).

The beneficial effect of CoQ10 on hematological parameters may be because of decreasing lipid peroxidation and inhibiting certain enzymes involved the formation of free radicals, blocking oxidative injuries to DNA and reducing glycation of membrane proteins (Modi et al. 2006, Prosek et al. 2008). Further, Ahmadvand et al. (2012) reported that CoQ10 reduced the lipid peroxidation and enhanced antioxidant enzyme activities (SOD, GSH, CAT) in experimentally diabetic rats. Quinzii et al. (2010) stated that there is strictly correlation reactive oxygen species, oxidative stress and cell death with CoQ10 deficiency.

On the other hand, experimentally induced diabetes lead to significantly (Table 2,  $p < 0.05$ ) increments in WBC count and neutrophil percentage compared to control group. Also, lymphocyte percentage was relatively decreased (Table 2,  $p < 0.05$ ) compared to control group. In diabetic rats treated





with CoQ10, neutrophil percentage and relative lymphocyte percentage were found to be lower and higher (Table 2,  $p < 0.05$ ) than that of diabetic group, respectively. In the same group, WBC count tended to decline but this changing was not important compared to diabetic group.

A positive association between increased levels of inflammatory markers (WBC count, CRP, and inflammatory cytokines) and diabetes incidence were determined in several studies (Gkrania-Klotsas et al. 2010, Twig et al. 2013). Tong et al. (2004) reported that elevated WBC count, even within the normal range, is associated with macro- and microvascular complications in diabetes. It has been noted that higher WBC count in diabetes reflects an inflammation and the other tissue complications (Tong et al. 2004, Twig et al. 2013). The increase neutrophil percentage by diabetes in this study is important regarding to show immune response against causal effects of diabetes. This result seems in coherent with above studies that noted about the same data obtained from diabetic subject. Although the decrease in WBC count is not important, significantly decrease in neutrophil percentage may be evaluated as a result of CoQ10 treatment's beneficial effect. There are many acknowledgements about anti-inflammatory properties of CoQ10. It was reported that CoQ10 administration in several doses decreased WBC count and inflammatory cytokines in human and animals (Ahmadvand et al. 2012, Abdollahzad et al. 2015). In the study conducted in animals, it has been suggested that CRP level were reduced with CoQ10 application (Devadasu et al. 2011). Above findings have supported our results obtained from CoQ10 administration in diabetic animals.

## Conclusion

In the light of our results, it can be said that CoQ10 supplementation has some ameliorative effects in respect to at least hematological parameters in streptozotocin-induced diabetic rats.

## References

- Abdollahzad H, Aghdashi MA, Asghari Jafarabadi M, Alipoor B, 2015. Effects of coenzyme Q10 supplementation on inflammatory cytokines (TNF- $\alpha$ , IL-6) and oxidative stress in rheumatoid arthritis patients: a randomized controlled trial. *Arch Med Res*, 46, 527-533.
- Ahmadvand H, Tavafi M, Khosrowbeygi A, 2012. Amelioration of altered antioxidant enzymes activity and glomerulosclerosis by coenzyme Q10 in alloxan-induced diabetic rats. *J Diabetes Complications*, 26, 476-482.
- Aladodo RA, Muhammad NO, Balogun EA, 2013. Effects of aqueous root extract of *Jatropha curcas* on hyperglycaemic and haematological indices in alloxan-induced diabetic rats. *FUJNAS*, 2, 52-58.
- American Diabetes Association, ADA, 2010. Diagnosis and classification of diabetes mellitus. *Diabetes care*, 33, 62-69.
- Bakan E, Yildirim A, Kurtul N, Polat MF, Dursun H, Cayir K, 2006. Effects of type 2 diabetes mellitus on plasma fatty acid composition and cholesterol content of erythrocyte and leukocyte membranes. *Acta Diabetol*, 43, 109-113.
- Brownlee M, 2001. Biochemistry and molecular cell biology of diabetic complications. *Nature*, 414, 813-820.
- Butler MG, Dasouki M, Bittel D, Hunter S, Naini A, DiMauro S, 2003. Coenzyme Q10 levels in Prader-Willi syndrome: comparison with obese and non-obese subjects. *Am J Med Genet A*, 119, 168-171.
- Cooke M, Iosia M, Buford T, Shelmadine B, Hudson G, Kerksick C, Rasmussen C, Greenwood M, Leutholtz B, Willoughby D, Kreider R, 2008. Effects of acute and 14-day coenzyme Q10 supplementation on exercise performance in both trained and untrained individuals. *J Int Soc Sports Nutr*, 5, 1-14.
- Crane FL, 2001. Biochemical functions of coenzyme Q10. *J Am Coll Nutr*, 20, 591-598.
- Devadasu VR, Wadsworth RM, Kumar MN, 2011. Protective effects of nanoparticulate coenzyme Q10 and curcumin on inflammatory markers and lipid metabolism in streptozotocin-induced diabetic rats: a possible remedy to diabetic complications. *Drug Deliv Transl Res*, 1, 448-455.
- Gkrania-Klotsas E, Ye Z, Cooper AJ, Sharp SJ, Luben R, Biggs ML, Chen LK, Gokulakrishnan K, Hanefeld M, Ingelsson E, Lai WA, Lin SY, Lind L, Lohsoonthorn V, Mohan V, Muscari A, Nilsson G, Ohrvik J, ChaoQiang J, Jenny NS, Tamakoshi K, Temelkova-Kurktschiev T, Wang YY, Yajnik CS, Zoli M, Khaw KT, Forouhi NG, Wareham NJ, Langenberg C, 2010. Differential white blood cell count and type 2 diabetes: systematic review and metaanalysis of cross-sectional and prospective studies. *PLoS ONE*, 5, e13405.
- Kennedy L, Baynes JW, 1984. Non-enzymatic glycosylation and the chronic complications of diabetes: an overview. *Diabetologia*, 26, 93-98.
- Labrousche S, Freyburger G, Gin H, Boisseau MR, Cassagne C, 1996. Changes in phospholipid composition of blood cell membranes (erythrocyte, platelet, and polymorphonuclear) in different types of diabetes clinical and biological correlations. *Metabolism*, 45, 57-71.
- Linnane AW, Kopsidas G, Zhang C, Yarovaya N, Kovalenko S, Papakostopoulos P, Eastwood H, Graves S, Richardson M, 2002. Cellular redox activity of coenzyme Q10: effect of CoQ10 supplementation on human skeletal muscle. *Free Radic Res*, 36, 445-453.
- Manodori AB, Kuypers FA, 2002. Altered red cell turnover in diabetic mice. *J Lab Clin Med*, 140, 161-165.
- Mansi KMS, 2006. Effects of administration of alpha-melanocyte stimulating hormone ( $\alpha$ -MSH) on some hematological values of alloxan-induced diabetic rats. *Am J Pharmacol Toxicol*, 1, 5-10.
- Modi K, Santani DD, Goyal RK, Bhatt PA, 2006. Effect of coenzyme Q10 on catalase activity and other antioxidant parameters in streptozotocin-induced diabetic rats. *Biol Trace Elem Res*, 109, 25-34.
- Moss PP, 1999. Blood banking: concepts and application. 12-34. P.A Saunders, Philadelphia.





- Muhammad NO, Oloyede OB, 2009. Haematological parameters of broiler chicks fed *Aspergillus niger*-fermented *Terminalia catappa* seed meal-based diet. *Global J Biotechnol Biochem*, 4, 179-183.
- Niki E, 1997. Mechanisms and dynamics of antioxidant action of ubiquinol. *Mol Aspects Med*, 18, 63-70.
- Niklowitz P, Menke T, Andler W, Okun JG, 2004. Simultaneous analysis of coenzyme Q10 in plasma, erythrocytes and platelets: comparison of the antioxidant level in blood cells and their environment in healthy children and after oral supplementation in adults. *Clin Chim Acta*, 342, 219-226.
- Prosek M, Butinar J, Lukanc B, Fir MM, Miliwojevic L, Krizman M, Smidovnik A, 2008. Bioavailability of water-soluble CoQ10 in beagle dogs. *J Pharm Biomed Anal*, 47, 918-922.
- Quinzii CM, Lopez LC, Gilkerson RW, Dorado B, Coku J, Naini AB, Lagier-Tourenne C, Schuelke M, Salviati L, Carrozzo R, Santorelli F, Rahman S, Tazir M, Koenig M, DiMauro S, Hirano M, 2010. Reactive oxygen species, oxidative stress, and cell death correlate with level of CoQ10 deficiency. *FASEB J*, 24, 3733-3743.
- Rosenfeldt F, Hilton D, Pepe S, Krum H, 2003. Systematic review of effect of coenzyme Q10 in physical exercise, hypertension and heart failure. *Biofactors*, 18, 91-100.
- Schmid-Schönbein H, Volger E, 1976. Red-cell aggregation and red cell deformability in diabetes. *Diabetes*, 25, 897-902.
- Sharma AK, 1993. *Diabetes mellitus and its complications: an update*. 92-205. 1st ed., New Delhi: Macmillan India Ltd.
- Shekelle P, Morton S, Hardy ML, 2003. Effect of supplemental antioxidants vitamin C, vitamin E, and coenzyme Q10 for the prevention and treatment of cardiovascular disease. *Evid Rep Technol Assess*, 83, 1-3.
- Tong PC, Lee KF, So WY, Ng MH, Chan WB, Lo MK, Chan NN, Chan JC, 2004. White blood cell count is associated with macro- and microvascular complications in Chinese patients with type 2 diabetes. *Diabetes Care*, 27, 216-222.
- Twig G, Afek A, Shamiss A, Derazne E, Tzur D, Gordon B, Tirosh A, 2013. White blood cells count and incidence of type 2 diabetes in young men. *Diabetes Care*, 36, 276-282.
- Wautier JL, Wautier MP, Schmidt AM, Anderson GM, Hori O, Zoukourian C, Capron L, Chappey O, Yan SD, Brett J, Guillausseau PJ, Stern D, 1994. Advanced glycation end products (AGEs) on the surface of diabetic erythrocytes bind to the vessel wall via a specific receptor inducing oxidant stress in the vasculature: a link between surface-associated AGEs and diabetic complications. *Proc Natl Acad Sci USA*, 91, 7742-7746.
- Zhou S, Zhang Y, Davie A, Marshall-Gradisnik S, Hu H, Wang J, Brushett D, 2005. Muscle and plasma coenzyme Q10 concentration, aerobic power and exercise economy of healthy men in response to four weeks of supplementation. *J Sports Med Phys Fitness*, 45, 337-346.

